arXiv:1504.00373v1 [math.RA] 1 Apr 2015

THE ASYMPTOTIC BEHAVIOR OF THE CODIMENSION SEQUENCE OF AFFINE G-GRADED ALGEBRAS

YUVAL SHPIGELMAN

Abstract. Let W be an affine PI algebra over a field of characteristic zero graded by a finite group G. We show that there exist $\alpha_1, \alpha_2 \in \mathbb{R}$, $\beta \in \frac{1}{2}\mathbb{Z}$, and $l \in \mathbb{N}$ such that $\alpha_1 n^\beta l^n \leq c_n^G(W) \leq \alpha_2 n^\beta l^n$. Furthermore, if W has a unit then the asymptotic behavior of $c_n^G(W)$ is $\alpha n^\beta l^n$ where $\alpha \in \mathbb{R}$, $\beta \in \frac{1}{2}\mathbb{Z}$, $l \in \mathbb{N}$.

Introduction

Throughout this article F is an algebraically closed field of characteristic zero and W is an affine associative F- algebra graded by a finite group G. We also assume that W is a PI-algebra; i.e., it satisfies an ordinary polynomial identity. In this article we study G-graded polynomial identities of W, and in particular the corresponding G-graded codimension sequence. Let us briefly recall the basic setup. Let X^G be a countable set of variables $\{x_{i,g} : g \in G; i \in \mathbb{N}\}$ and let $F\langle X^G\rangle$ be the free algebra on the set X^G. Given a polynomial in $F\langle X^G\rangle$ we say that it is a G-graded identity of W if it vanishes upon any admissible evaluation on W. That is, a variable $x_{i,g}$ assumes values only from the corresponding homogeneous component W_g. The set of all G-graded identities is an ideal in the free algebra $F\langle X^G\rangle$ which we denote by $Id^G(W)$. Moreover, $Id^G(W)$ is a G-graded T-ideal, namely, it is closed under G-graded endomorphisms of $F\langle X^G\rangle$. Let $R^G(W)$ denote the relatively free algebra $F\langle X^G\rangle / Id^G(W)$, and $C_n^G(W)$ denote the space $P_n^G / P_n^G \cap Id^G(A)$ where P_n^G is the $|G|^n \cdot n!$ dimensional F-space spanned by all permutations of the (multilinear) monomials $x_{1,g_1}, x_{2,g_2} \cdots x_{n,g_n}$ where $g_i \in G$. We also define the n-th coefficient $c_n^G(W)$ of the codimension sequence of W by $c_n^G(W) = dim_F\left(C_n^G(W)\right)$. In [AB2] Aljadeff and Belov showed that for every affine G-graded algebra there exists a finite dimensional G-graded algebra with the same ideal of graded identities, thus with the same codimension sequence. We denote such algebra by A_W.

In the last few years several papers have been written which generalize results from the theory of ordinary polynomial identities to the G -graded case (e.g. see [Sv, AB1, AB2]). One of these papers was by E.Aljadeff, A.Giambruno and D.La Matina who showed, in different collaborations (see [GL, AG, AGL]), that, as in the nongraded case, $\lim_{n\to\infty} \sqrt[n]{c_n^G(W)}$ exists, and it is a nonnegative integer called the G - graded exponent of W, and denoted by $exp^G(W)$. In this paper we expand this result and prove that,

Key words and phrases. graded algebras, polynomial identities, representation theory, Hilbert series, codimension.

in addition to the "exponential part", there is a "polynomial part" to the asymptotics of $c_n^G(W)$. More precisely, we prove :

Theorem (***A***)**.** *Let G be a finite group, and W an affine G- graded F-algebra where F is a field of characteristic 0. Suppose that W satisfies an ordinary polynomial identity. Then there exist $\alpha_1, \alpha_2 > 0$, $\beta \in \frac{1}{2}\mathbb{Z}$, and $l \in \mathbb{N}$ such that*

$$\alpha_1 n^\beta l^n \leq c_n^G(W) \leq \alpha_2 n^\beta l^n.$$

A conclusion of this theorem is that $\lim_{n\to\infty} log_n\left(\frac{c_n(W)}{exp(W)^n}\right)$ (the power of the "polynomial part") exists, and is an integer or a half-integer.

Furthermore, if W has a unit, we have found the structure of the codimension sequence's asymptotics.

Theorem (**B**)**.** *Let G be a finite group, and W an unitary affine G- graded F-algebra where F is a field of characteristic 0. Suppose that W satisfies an ordinary polynomial identity. Then there exist $\beta \in \frac{1}{2}\mathbb{Z}$, $l \in \mathbb{N}$, and $\alpha \in \mathbb{R}$ such that*

$$c_n^G(W) \sim \alpha n^\beta l^n.$$

Theorems ***A*** and **B** generalize results from the theory of ordinary polynomial identities. Indeed, in [BR] Berele and Regev proved the nongraded version of theorems ***A*** and **B** for PI algebras satisfying a Capelli identity (e.g affine algebras, see [GZ2]). However, in Theorem A (for ungraded algebras) Berele and Regev made an additional assumption, namely, that the codimension sequence of the PI algebra is eventually nondecreasing. Recently, in [GZ3], Giambruno and Zaicev showed that the codimension sequence of any PI algebra is eventually nondecreasing and therefore the additional assumption mentioned above can be removed. Here we prove a G-graded version of Giambruno and Zaicev's result and so the assumption on the monotonicity of the G-greded codimension sequence is also unnecessary.

We begin our article (Section 1) by mentioning some notations from the representation theory of $GL_n(F)$ and S_n. Section 2 is dealing with the case of affine (not necessarily unitary) G - graded algebras (Theorem A). And Section 3 is about the case of unitary affine G-graded algebras (Theorem B).

1. Preliminaries

In this section we recall some notations and definitions from the representation theory of $GL_n(F)$ and S_n.

Definition 1.1. A *partition* is a finite sequence of integers $\lambda = (\lambda_1, ..., \lambda_k)$ such that $\lambda_1 \geq \cdots \geq \lambda_k > 0$. The integer k denoted by $h(\lambda)$, and called the *height* of λ. The set of all partitions is denoted by Λ, and the set of all partitions of height less or equal to k is denoted by Λ^k .

We say that λ is a *partition of* $n \in \mathbb{N}$ if $\sum_{i=1}^{k} \lambda_i = n$. In this case we write $\lambda \vdash n$ or $|\lambda| = n$.

It is known (e.g. see [Bru]) that the irreducible S_n representations are indexed by partitions of n. Denote by D_λ the S_n irreducible representation indexed by λ, and by χ_λ the corresponding character. By Maschke's theorem every S_n - representation V is completely reducible, so we can write $V = \bigoplus_{\lambda \vdash n} m_\lambda D_\lambda$, and its character

$$\chi(V) = \sum_{\lambda \vdash n} m_\lambda \chi_\lambda.$$

A $GL_n(F)$ - representation $Y = Sp_F\{y_1, y_2, ...\}$ is called a *polynomial* if there is a set of polynomials $\{f_{i,j}(z_{s,t}) | i, j \in \mathbb{N}\} \subset F[z_{s,t} | 1 \leq s, t \leq n]$ such that for every i only finite number of $f_{i,j}(z_{s,t})$'s are nonzero, and the action of $GL_n(F)$ on Y is given by:

$$P \cdot y_i = \sum_j f_{i,j}(p_{s,t}) y_j$$

for every $P = (p_{s,t}) \in GL_n(F)$. We say that Y is $q-$*homogeneous* if all the nonzero $f_{i,j}$'s are homogeneous polynomials of (total) degree q. For $\alpha = (\alpha_1, ..., \alpha_n) \in \mathbb{N}^n$, one defines the *weight space* of Y associated to α by

$$Y^\alpha = \{y \in Y | \mathrm{diag}(z_1, ..., z_n) \cdot y = z_1^{\alpha_1} \cdots z_n^{\alpha_n} y\}.$$

It is known that

$$Y = \bigoplus_\alpha Y^\alpha.$$

The series $H_Y(t_1, ..., t_n) = \sum_\alpha (\dim_F Y^\alpha) t_1^{\alpha_1} \cdots t_n^{\alpha_n}$ is called the *Hilbert (or Poincare) series* of Y. The Hilbert series is known to be symmetric in $t_1, ..., t_n$, so we recall an important basis of the space of symmetric series, namely the Schur functions. For convenience we use the following combinatorial definition: (Note that although this definition is not the classical one , it is equivalent to it e.g. see [Bru])

Let $\lambda = (\lambda_1, .., \lambda_k)$ be a partition. The *Young diagram* associated with λ is the finite subset of $\mathbb{Z} \times \mathbb{Z}$ defined as $\mathcal{D}_\lambda = \{(i, j) \in \mathbb{Z} \times \mathbb{Z} | i = 1, .., k\,,\ j = 1, .., \lambda_i\}$. We may regard $\mathcal{D}_\lambda$ as k arrays of boxes where the top one is of length λ_1, the second of length λ_2, etc. For example

$\mathcal{D}_{(4,3,3,1)} =$ 

A Schur function $s_\lambda \in \mathbb{Z}[t_1, ..., t_n]^{S_n}$ is a polynomial such that the coefficient of $t_1^{a_1} \cdots t_n^{a_n}$ is equal to the number of ways to insert a_1 ones, a_2 twos, ... , and a_n n's in $\mathcal{D}_\lambda$ such that in every row the numbers are non-decreasing, and in any column the numbers are strictly increasing. Note that s_λ is homogenous of degree $|\lambda|$.

Example.

(1) If λ is a partition of height one, i.e $\lambda = (\lambda_1)$, then the corresponding Schur function is

$$s_{(\lambda_1)}(t_1, ..., t_n) = \sum_{a_1+\cdots+a_n=\lambda_1} t_1^{a_1} \cdots t_n^{a_n}.$$

(2) If $\lambda = (2, 1)$ and $n = 2$ we have

$$s_{(2,1)}(t_1, t_2) = t_1^2 t_2 + t_1 t_2^2$$

since the only two ways to set ones and twos in $\mathcal{D}_{(2,1)}$ is $\begin{array}{|c|c|}\hline 1 & 1 \\ \hline 2 \\ \cline{1-1}\end{array}$ and $\begin{array}{|c|c|}\hline 1 & 2 \\ \hline 2 \\ \cline{1-1}\end{array}$.

The next lemma is easily implied from the definition of the Schur functions.

Lemma 1.2.

(1) $s_\lambda(t_1, ..., t_n) = 0$ *if and only if* $h(\lambda) > n$.
(2) $s_\lambda(t_1, ..., t_n, 0, ..., 0) = s_\lambda(t_1, ..., t_n)$.

As $\{s_\lambda(t_1, ..., t_n)\}_{\lambda\in\Lambda^n}$ inform a basis of the space of symmetric series in n variables there exist integers $\{m_\lambda\}_{\lambda\in\Lambda^n}$ such that

$$H_Y(t_1, ..., t_n) = \sum_{\lambda\in\Lambda^n} m_\lambda s_\lambda(t_1, ..., t_n)$$

On the other hand, the group S_n is embed in $GL_n(F)$ via permutation matrices, hence Y is also an S_n representation. Note that if P is a permutation matrix there is $\sigma \in S_n$ such that $\mathrm{diag}(z_1, ..., z_n) \cdot P = P \cdot \mathrm{diag}(z_{\sigma(1)}, ..., z_{\sigma(n)})$. Hence for every y in the weight space $Y^{(1^n)}$ (1^n means $1, ..., 1$ n times) we have

$$\mathrm{diag}(z_1, ..., z_n) \cdot Py = P \cdot \mathrm{diag}(z_{\sigma(1)}, ..., z_{\sigma(n)})y = z_1 \cdots z_n Py$$

so $Py \in Y^{(1^n)}$, and we obtain that $Y^{(1^n)}$ is a sub S_n representation of Y. Moreover, by the representation theory of $GL_n(F)$ the S_n character of $Y^{(1^n)}$ is $\sum_{\lambda\vdash n} m_\lambda \chi_\lambda$ with the same coefficients m_λ as in the Hilbert series of Y.

Next we recall some terminology and facts about G - graded relatively free algebras. Denote $X_n^G = \{x_{i,g} | g \in G\,,\, 1 \le i \le n\} \subset X^G$, and consider the free algebra $F\left\langle X_n^G\right\rangle$ which is a subalgebra of $F\left\langle X^G\right\rangle$. We equip $F\left\langle X_n^G\right\rangle$ with a G-grading by setting the homogenous degree of a monomial $x_{i_1,g_{i_1}} \cdots x_{i_l,g_{i_l}}$ to be $g_{i_1} \cdots g_{i_l} \in G$. If we suppose that $\mathcal{I}$ is a G - graded T-ideal of the free algebra $F\left\langle X_n^G\right\rangle$, then it is easy to see that the grading on $F\left\langle X_n^G\right\rangle$ induces (naturally) a grading on ${}^{F\left\langle X_n^G\right\rangle}/_{\mathcal{I}}$, giving us the corresponding T-ideal of G-graded identities $Id^G\left({}^{F\left\langle X_n^G\right\rangle}/_{\mathcal{I}}\right) \subset F\left\langle X^G\right\rangle$.
We say that $\mathcal{I}$ is PI if $Id^G\left({}^{F\left\langle X_n^G\right\rangle}/_{\mathcal{I}}\right) \neq \{0\}$ i.e.${}^{F\left\langle X_n^G\right\rangle}/_{\mathcal{I}}$ is a PI algebra. Note that the T-ideal $F\left\langle X_n^G\right\rangle \cap Id^G\left({}^{F\left\langle X_n^G\right\rangle}/_{\mathcal{I}}\right)$ is exactly $\mathcal{I}$.

For every $g \in G$ the group $GL_n(F)$ acts linearly on the n-dimensional vector space $Sp_F\{x_{1,g}, ..., x_{n,g}\}$. Given these actions, we define a polynomial $GL_n(F)$ representation

structure on $F\left\langle X_n^G\right\rangle$ via $P(x_{i_1,g_1}\cdots x_{i_l,g_l})=P(x_{i_1,g_1})\cdots P(x_{i_l,g_l})$ for every $P\in GL_n(F)$. This structure induces (naturally) a structure of $GL_n(F)$ representation on the relatively free algebra ${}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{I}}$, giving us the corresponding Hilbert series $H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{I}}}(t_1,...,t_n)$. We recall from [AB1] (lemma 2.10 and 2.11) the following properties of that kind of Hilbert series:

Lemma 1.3. *Let $\mathcal{J}_1,\mathcal{J}_2$ be G - graded T-ideals of $F\left\langle X_n^G\right\rangle$, then*

(1) $H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{J}_1\cap\mathcal{J}_2}}=H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{J}_1}}+H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{J}_2}}-H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{J}_1+\mathcal{J}_2}}$.
(2) *If $\mathcal{J}_1\subset\mathcal{J}_2$ then*

$$H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{J}_1}}=H_{{}^{F\left\langle X_n^G\right\rangle}\!/_{\mathcal{J}_2}}+H_{\mathcal{J}_2/\mathcal{J}_1}.$$

In our case recall that $Id_n^G(W):=F\left\langle X_n^G\right\rangle\cap Id^G(W)$ is G-graded T-ideal which is PI for every $n\in\mathbb{N}$, and let

$$\left\{R_n^G(W)={}^{F\left\langle X_n^G\right\rangle}\!/_{Id_n^G(W)}\right\}_{n\in\mathbb{N}}$$

be a collection of affine relatively free G-graded algebras of W, and consider the corresponding Hilbert series $H_{R_n^G(W)}(t_1,...,t_n)=\sum_\lambda m_\lambda^{(n)}s_\lambda(t_1,...,t_n)$. We can show that we need only one set of coefficients $\{m_\lambda\}_{\lambda\in\Lambda}$ to describe $H_{R_n^G(W)}(t_1,...,t_n)$ for every n. Indeed, it is easy to see that by definition $\mathrm{diag}(z_1,...,z_n)\cdot x_{i_1,g_1}\cdots x_{i_l,g_l}=z_{i_1}\cdots z_{i_l}x_{i_1,g_1}\cdots x_{i_l,g_l}$, hence the weight space $\left(R_n^G(W)\right)^\alpha$ is the span of all the monomials in $R_n^G(W)$ with α_i appearances of variables from the set $\{x_{i,g}|g\in G\}$ for every $i=1,...,n$. Therefore, we can write $R_r^G(W)=\bigoplus\left(R_m^G(W)\right)^\alpha$ for every $r<m$, where the summing is over all the $\alpha\in\mathbb{N}^m$ such that $\alpha=(\alpha_1,...,\alpha_r,0,...,0)$. Thus, we get the Hilbert series of $R_r^G(W)$ by putting $t_{r+1}=\cdots=t_m=0$ in the Hilbert series of $R_m^G(W)$, i.e.

$$H_{R_r^G(W)}(t_1,...,t_r)=\sum_{\lambda\in\Lambda^r}m_\lambda^{(r)}s_\lambda(t_1,...,t_r)=\sum_{\lambda\in\Lambda^m}m_\lambda^{(m)}s_\lambda(t_1,...,t_r,0,...0).$$

Moreover, by Lemma 1.2 we get

$$\sum_{\lambda\in\Lambda^r}m_\lambda^{(r)}s_\lambda(t_1,...,t_r)=\sum_{\lambda\in\Lambda^r}m_\lambda^{(m)}s_\lambda(t_1,...,t_r)+\sum_{\lambda\in\Lambda^m\setminus\Lambda^r}m_\lambda^{(m)}\cdot 0,$$

hence for every $r<m$ the coefficients $m_\lambda^{(r)}$ and $m_\lambda^{(m)}$ are the same for every $\lambda\in\Lambda^r$. Making it possible to denote $m_\lambda^{(n)}=m_\lambda$, and write for every $n\in\mathbb{N}$

$$H_{R_n^G(W)}(t_1,...,t_n)=\sum_{\lambda\in\Lambda^n}m_\lambda s_\lambda(t_1,...,t_n).$$

Remember that A_W is a finite dimensional G-graded algebra such that $Id^G(A_W)=Id^G(W)$, thus with the same m_λ. By [Go] (Lemma 1 and Lemma 7) $m_\lambda=0$ if $\lambda\notin\Lambda^k$ where $k=dim_F(A_W)$, so all the nonzero coefficients m_λ appear in the series $H_{R_k^G(W)}(t_1,...,t_k)$. Notice that for every n the weight space $R_n^G(W)^{(1^n)}$ is exactly $C_n^G(W)$, thus $c_n^G(W)=\sum_{\lambda\vdash n}m_\lambda d_\lambda$ where $d_\lambda=dim_F D_\lambda$. The conclusion is that the G - graded

codimension sequence of W is determined by a single Hilbert series $H_{R_k^G(W)}(t_1,...,t_k)$. We denote this series by $H_W(t_1,...,t_k)$.

As mention in the introduction, Berele and Regev proved versions of Theorems ***A*** and **B** for (nongraded) affine PI algebras. Their proofs rely on a result about symmetric series which we will also use. Before stating this result we need a definition.

Definition 1.4. A series $H(t_1,...,t_k)$ is called *rational function* if we can write it as $\frac{f(t_1,...,t_k)}{g(t_1,...,t_k)}$ where f,g are polynomials, moreover a rational function $\frac{f(t_1,...,t_k)}{g(t_1,...,t_k)}$ is called *nice* if $g(t_1,...,t_k)=\prod_{\alpha\in\Delta}(1-t_1^{\alpha_1}\cdots t_k^{\alpha_k})$ where Δ is some subset of $\mathbb{N}^k$.

Theorem 1.5 ([BR]). *Let $H(t_1,...,t_k)=\sum_{\lambda\in\Lambda}a_\lambda s_\lambda(t_1,...,t_k)$ be a series, and denote $y_n^H=\sum_{\lambda\vdash n}a_\lambda d_\lambda$ where d_λ is the dimension of the λ's irreducible representation of S_n, and $a_\lambda\in\mathbb{N}$. We assume that*

1. *There is an integer r such that $a_\lambda=0$ for every partition λ satisfying $\lambda_{r+1}\geq M$ where M is a constant which depends only on r.*
2. *$H(t_1,...,t_k)$ is nice rational function.*

Then there is an integer d such that for every $0\leq m\leq d-1$ there are constants $\beta_m\in\frac{1}{2}\mathbb{Z}$, and $\alpha_m\geq 0$. such that

$$y^H_{n_{m,t}}\sim\alpha_m n_{m,t}^{\beta_m}l^{n_{m,t}}\quad;\quad t\to\infty$$

where $n_{m,t}=m+td$, and l is the minimal integer satisfying condition 1.

2. Affine G-graded Algebras

In this section we prove Theorem A. Our first step is to show that $H_W(t_1,...,t_k)=\sum_{\lambda\in\Lambda^k}m_\lambda s_\lambda$ satisfies the conditions of Theorem 1.5, hence $y_n^{H_W}=c_n^G(W)$ satisfies the conclusion of the theorem. Recall that according to the first condition there is an integer r such that $m_\lambda=0$ for every partition λ satisfying $\lambda_{r+1}\geq M$ where M is a constant which depends only on r. In [Go] A.S Gordienko (see Lemma 1 and Lemma 7) showed that this condition holds for finite dimensional G-graded PI F-algebras, however we know (see the introduction) that there is a finite dimensional G-graded algebra A_W such that $H_W=H_{R_k^G(A_W)}$, so indeed H_W satisfies the first condition of Theorem 1.5. Let us show that the second condition of Theorem 1.5 holds, namely that H_W is nice rational. In [AB1] Aljadeff and Belov proved that the Hilbert series of ${}^{F\langle X_k^G\rangle}/_{\mathcal{I}}$ is rational for every G - graded T - ideal $\mathcal{I}$ of $F\left\langle X_k^G\right\rangle$ which is PI. In fact, with slight changes in their proof, we can prove that the Hilbert series is also nice.

Theorem 2.1. *Let $\mathcal{I}$ be a G - graded T - ideal of $F\left\langle X_k^G\right\rangle$ which is PI then the Hilbert series of ${}^{F\langle X_k^G\rangle}/_{\mathcal{I}}$ is a nice rational function.*

Proof. Assume by contradiction that $H_{{}^{F\langle X_k^G\rangle}/_{\mathcal{I}}}(t_1,...,t_k)$ is not nice rational. Then the set of all PI G - graded T-ideal of $F\left\langle X_k^G\right\rangle$ whose Hilbert series is not nice rational, is

not empty. Furthermore, we may assume that there exists a maximal element in this set, because otherwise (by Zorn's lemma) there is an infinite ascending sequence in this set. This sequence does not stabilize, and hence its union is a nonfinite generated T-ideal. This of course contradicts the Specht property of G- graded T-ideals (see [AB2] section 12).

So let $\mathcal{J}$ be a maximal PI G-graded T-ideal of $F\left\langle X_k^G\right\rangle$ such that $H_{F\left\langle X_k^G\right\rangle/\mathcal{J}}$ is not nice rational. From [AB2] (Theorem 11.2) we know that

$$Id^G\left(F\left\langle X_k^G\right\rangle/\mathcal{J}\right) = Id^G(A_1 \oplus \cdots \oplus A_l) \tag{2.1}$$

where A_i is a G-graded basic F-algebra for every i, and $id^G(A_i) \nsubseteq id^G(A_j)$ for any $1 \leq i, j \leq m$ with $i \neq j$. We refer the reader to [AB2] for more details - in particular, the definition of basic algebra. For our purpose, we just have to use one property of basic algebras which we will present later. Now, from 2.1 we get that $\mathcal{J} = Id_k^G\left(F\left\langle X_k^G\right\rangle/\mathcal{J}\right) = Id_k^G(A_1) \cap Id_k^G(A_2 \oplus \cdots \oplus A_l)$. Therefore, if we denote $\mathcal{J}_1 = Id_k^G(A_1)$ and $\mathcal{J}_2 = Id_k^G(A_2 \oplus \cdots \oplus A_l)$ then we get from Lemma 1.3

$$H_{F\left\langle X_k^G\right\rangle/\mathcal{J}} = H_{F\left\langle X_k^G\right\rangle/\mathcal{J}_1} + H_{F\left\langle X_k^G\right\rangle/\mathcal{J}_2} - H_{F\left\langle X_k^G\right\rangle/\mathcal{J}_1+\mathcal{J}_2}.$$

Now, if $l > 1$ then $\mathcal{J} \subsetneq \mathcal{J}_1, \mathcal{J}_2$ from the maximality of $\mathcal{J}$ the series $H_{F\left\langle X_k^G\right\rangle/\mathcal{J}_1}$, $H_{F\left\langle X_k^G\right\rangle/\mathcal{J}_2}$ and $H_{F\left\langle X_k^G\right\rangle/\mathcal{J}_1+\mathcal{J}_2}$ are all nice rational functions. Therefore, $H_{F\left\langle X_k^G\right\rangle/\mathcal{J}}$ is nice rational which is a contradiction. Hence, $l = 1$, and

$$Id^G\left(F\left\langle X_k^G\right\rangle/\mathcal{J}\right) = Id^G(A)$$

where A is a basic F-algebra.

We recall from [AB1] (see section 2) that for any basic algebra A there is a G-graded T-ideal $\mathcal{K}$ strictly containing $\mathcal{J} = Id_k^G(A)$ such that $\mathcal{K}/\mathcal{J}$ is a finite module over some affine commutative F-algebra, hence by [St] (Chapter 1 Theorem 2.3) $H(\mathcal{K}/\mathcal{J})$ is a nice rational function. Now, by the maximality of $\mathcal{J}$ the Hilbert series of $F\left\langle X_k^G\right\rangle/\mathcal{K}$ is nice rational. Moreover, by Lemma 1.3 we get

$$H_{F\left\langle X_k^G\right\rangle/\mathcal{J}} = H_{F\left\langle X_k^G\right\rangle/\mathcal{K}} + H_{\mathcal{K}/\mathcal{J}}$$

hence $H_{F\left\langle X_k^G\right\rangle/\mathcal{J}}$ is nice rational which contradicts our assumption. □

So indeed we can apply Theorem 1.5 on H_W, and we arrive at the following corollary for affine G-graded algebra with eventually non-decreasing codimension sequence (that is $c_n^G(W) \leq c_{n+1}^G(W)$ for every $n > N$ for some $N \in \mathbb{N}$).

Corollary 2.2. *If the sequence $c_n^G(W)$ is eventually non-decreasing then there exists $\alpha_1, \alpha_2 > 0$, $\beta \in \frac{1}{2}\mathbb{Z}$, and $l \in \mathbb{N}$ such that*

$$\alpha_1 n^\beta l^n \leq c_n^G(W) \leq \alpha_2 n^\beta l^n.$$

Proof. We need to show that all the β_m's form Theorem 1.5 are equal. Suppose that there are $0 \leq m, l \leq d-1$ such that $\beta_m < \beta_l$, and denote $s = l - m \pmod{d}$. Note that $s = m - l + qd$ thus

$$n_{m,t} + s = m + td + s = m + td + l - m + qd = l + (t+q)d = n_{l,t+q}$$

therefore

$$\lim_{t\to\infty} \frac{c^G_{n_{m,t}+s}(W)}{c^G_{n_{m,t}}(W)} = \lim_{t\to\infty} \frac{c^G_{n_{l,t+q}}(W)}{c^G_{n_{m,t}}(W)} = \lim_{t\to\infty} \frac{\alpha_l {n_{l,t}}^{\beta_l} l^{n_{l,t}}}{\alpha_m n_{m,t}^{\beta_m} l^{n_{m,t}}} = 0$$

which is a contradiction since $c^G_n(W)$ is eventually non-decreasing. □

The final stage of Theorem A's proof is removing the non-decreasing assumption from 2.2.

Proposition 2.3. *Let G be a finite group, and W an affine G- graded F-algebra where F is a field of characteristic 0. Suppose that W satisfying an ordinary polynomial identity. Then the G- graded codimension sequence of W is eventually non-decreasing.*

Proof. Recall that there exists a finite dimensional algebra A_W having the same codimension sequence as W, so we may prove the proposition for A_W. Let $A_W = S + J$ be the Wedderburn-Malcev decomposition of A_W, and we may assume that the semi-simple part S, and the Jacobson radical J are G-graded. It is known (see [BRo]) that J is nilpotent, i.e. there is an integer t such that $J^t = 0$. We claim that $c^G_n(A_W) \leq c^G_{n+1}(A_W)$ for every $n > t$.

If $S = 0$ then A_W is nilponent of degree t, thus $c^G_n(A_W) = 0$ for every $n > t$. So assume that $S \neq 0$, and let 1_S be its unit element 1_S. Because $1_S^2 = 1_S$, the linear operator $T : A_W \to A_W$; $T(x) = x \cdot 1_S$ has eigenvalues 1 or 0, so we have the decomposition $A_W = A_0 \oplus A_1$ where A_0, A_1 are the eigenspaces of 0 and 1 respectively. Now, let $B = B_0 \cup B_1$ be a basis of A_W where $B_0 \subset A_0$ and $B_1 \subset A_1$. We may assume that all the elements in the basis are G- homogenous. Since F is of characteristic 0, it is enough to substitute the variables in a polynomial $f \in F\langle X^G \rangle$ by element form B in order to determine if f is an identity or not.

Denote $G = \{g_1, ..., g_s = e\}$, and for any $\mathbf{h} = (h_1, ..., h_n) \in G^n$ denote $P_{\mathbf{h}} = Sp_F\{x_{1,h_1}, ..., x_{n,h_n}\}$ and $C_{\mathbf{h}} = {}^{P_{\mathbf{h}}}/_{P_{\mathbf{h}} \cap Id^G(A_W)}$. Note that $C^G_n(A_W) = \bigoplus_{\mathbf{h}\in G^n} C_{\mathbf{h}}$, and $C_{\mathbf{h}} \cong C_{\mathbf{k}}$ if the vector $\mathbf{k}$ is a permutation of the vector $\mathbf{h}$. Therefore, if we denote $P_{(g_1^{n_1},...,g_s^{n_s})} = P_{n_1,...,n_s}$ and $C_{(g_1^{n_1},...,g_s^{n_s})} = C_{n_1,...,n_s}$ ($g_i^{n_i}$ means $g_i, ..., g_i$ n_i times), we can write

$$c^G_n(A_W) = \sum_{n_1+\cdots+n_s=n} \binom{n}{n_1, ..., n_s} c_{n_1,...,n_s} \tag{2.2}$$

where $\binom{n}{n_1,...,n_s} = \frac{n!}{n_1!\cdots n_s!}$ is the generalized binomial coefficient and

$$c_{n_1,...,n_s} = dim_F C_{n_1,...,n_s}.$$

Given $t < n = n_1 + \cdots + n_s$, write $c_{n_1,\ldots,n_s} = k$ and let $f_1(x_{1,h_1}, \ldots, x_{n,h_n}), \ldots, f_k(x_{1,h_1}, \ldots, x_{n,h_n})$ be a set of linear independent polynomials in ${}^{P_{n_1,\ldots,n_s}}/_{P_{n_1,\ldots,n_s} \cap Id^G(W)}$ (note that $h_1 = \cdots = h_{n_1} = g_1$, $h_{n_1+1} = \cdots = h_{n_1+n_2} = g_2$ and so on). For every $i = 1, \ldots k$ we define the following polynomials in $P_{n_1,\ldots,n_s+1}$ (recall that $g_s = e$, so $P_{n_1,\ldots,n_s+1} = P_{(h_1,\ldots,h_n,e)}$).

$$p_i(x_{1,h_1}, \ldots, x_{n,h_n}, x_{n+1,e}) = \sum_{j=1}^{k} f_i(x_{1,h_1}, \ldots, x_{j,h_j} x_{n+1,e}, \ldots, x_{n,h_n}).$$

We want to show that $p_1, \ldots, p_k \in {}^{P_{n_1,\ldots,n_s+1}}/_{P_{n_1,\ldots,n_s+1} \cap Id^G(W)}$ are linearly independent, so let $\sum_{i=1}^{k} \alpha_i p_i$ be a nontrivial linear combination of the p_i's. Since $f_1, \ldots, f_k$ are linear independent, there are $b_{j,h_j} \in B \cap (A_W)_{h_j}$ such that $\sum_{i=1}^{k} \alpha_i f_i(b_{1,h_1}, \ldots, b_{n,h_n}) \neq 0$. Since n is greater than the nilpotency index of J, one of the b_{i,h_i}'s is necessarily in S, and $S \subset A_1$ so $q = |B_1 \cap \{b_{1,h_1}, \ldots, b_{n,h_n}\}|$ is a nonzero integer. Let us substitute $b_{1,h_1}, \ldots, b_{n,h_n}, 1_S$ in p_i. Note that $b_{j,h_j} \cdot 1_S = b_{j,h_j}$ if $b_{j,h_j} \in B_1$, and $b_{j,h_j} \cdot 1_S = 0$ if $b_{j,h_j} \in B_0$, therefore

$$p_i(b_{1,h_1}, \ldots, b_{n,h_n}, 1_S) = q \cdot f_i(b_{1,h_1}, \ldots, b_{n,h_n})$$

and

$$\sum_{i=1}^{k} \alpha_i p_i(b_{1,h_1}, \ldots, b_{n,h_n}, 1_S) = q \cdot \sum_{i=1}^{k} \alpha_i f_i(b_{1,h_1}, \ldots, b_{n,h_n}) \neq 0.$$

Thus the polynomial $\sum_{i=1}^{k} \alpha_i p_i(x_{1,h_1}, \ldots, x_{n,h_n}, x_{n+1,e})$ is not an identity, so

$$c_{n_1,\ldots,n_s} = k \leq c^G_{n_1,\ldots,n_s+1}$$

for every $n_1, \ldots, n_s$. Hence by 2.2

$$c^G_n(W) = \sum_{n_1+\cdots+n_s=n} \binom{n}{n_1, \ldots, n_s} c_{n_1,\ldots,n_s}$$

$$\leq \sum_{n_1+\cdots+n_s=n} \binom{n+1}{n_1, \ldots, n_s+1} c^G_{n_1,\ldots,n_s+1} \leq c^G_{n+1}(W)$$

where $\binom{n}{n_1,\ldots,n_s} \leq \binom{n+1}{n_1,\ldots,n_s+1}$ is due to the combinatorial identity

$$\binom{n+1}{n_1, \ldots, n_s+1} = \binom{n}{n_1, \ldots, n_s} + \sum_{i=1}^{s-1} \binom{n}{n_1, \ldots, n_i - 1, \ldots, n_s}.$$

□

From 2.2 and 2.3 we conclude:

Theorem (***A***)**.** *Let G be a finite group, and W an affine G- graded F-algebra where F is a field of characteristic 0. Suppose W satisfies an ordinary polynomial identity.*

Then there exists $\alpha_1, \alpha_2 > 0$, $\beta \in \frac{1}{2}\mathbb{Z}$, *and* $l \in \mathbb{N}$ *such that*

$$\alpha_1 n^\beta l^n \leq c_n^G(W) \leq \alpha_2 n^\beta l^n.$$

A conclusion drawn from theorem A is that the codimension sequence's asymptotics has a "polynomial part" of degree $\beta \in \frac{1}{2}\mathbb{Z}$. More precisely,

Corollary 2.4. *The following limit exists and it is a half integer.*

$$\lim_{n\to\infty} log_n \left(\frac{c_n^G(W)}{exp^G(W)^n} \right)$$

3. Unitary Affine G-graded Algebras

In this section we assume that W has a unit. We show that, as in the nongraded case, the asymptotics of the codimension sequence of W is a constant times polynomial part times exponential part (Theorem B).

In [D] V.Drensky showed that in the non-graded case (or in another words for $G = \{e\}$) there is a subspace B_n of $R_n^{\{e\}}(W)$ such that $R_n^{\{e\}}(W) = F[z_1, ..., z_n] \otimes B_n$ for every $n \in \mathbb{N}$. We start this section by proving this result for any finite group G. In order to keep the notations as light as possible, whenever there is an element $f + I$ in some quotient space A/I where $A \subset F\langle X_n^G \rangle$ we omit the I and leave only the polynomial f when the convention is that all the operations on such polynomials are modulo I. So let B_n be the subspace of $R_n^G(W)$ spanned by all the polynomials $x_{i_1,h_1} \cdots x_{i_r,h_r}[x_{i_{r+1},g_1}, ..., x_{i_{r+l},g_l}] \cdots [..., x_{i_s,g_s}]$ where $h_j \neq e$ for $i = 1, ..., r$. Note that B_n is clearly a sub polynomial $GL_n(F)$ representation of $R_n^G(W)$, so the Hilbert series $H_{B_n}(t_1, ..., t_n) = \sum_{\lambda\in\Lambda^n} a_\lambda^{(n)} s_\lambda(t_1, ..., t_n)$ is well defined. Moreover, we can write $B_r = \bigoplus B_m^\alpha$ for every $r < m$, where the sum is over all the $\alpha \in \mathbb{N}^m$ such that $\alpha = (\alpha_1, ..., \alpha_r, 0, ..., 0)$. Thus, we can get the Hilbert series of B_r by putting $t_{r+1} = \cdots = t_m = 0$ in the Hilbert series of B_m, i.e.

$$H_{B_r}(t_1, ..., t_r) = \sum_{\lambda\in\Lambda^r} a_\lambda^{(r)} s_\lambda(t_1, ..., t_r) = \sum_{\lambda\in\Lambda^m} a_\lambda^{(m)} s_\lambda(t_1, ..., t_r, 0, ...0).$$

Moreover, from Lemma 1.2

$$\sum_{\lambda\in\Lambda^r} a_\lambda^{(r)} s_\lambda(t_1, ..., t_r) = \sum_{\lambda\in\Lambda^r} a_\lambda^{(m)} s_\lambda(t_1, ..., t_r) + \sum_{\lambda\in\Lambda^m\setminus\Lambda^r} a_\lambda^{(m)} \cdot 0.$$

Hence, for every $r < m$ the coefficients $a_\lambda^{(r)}$ and $a_\lambda^{(m)}$ are the same for every $\lambda \in \Lambda^r$. Making it possible to denote $a_\lambda^{(n)} = a_\lambda$, and write for every $n \in \mathbb{N}$

$$H_{B_n} = \sum_{\lambda\in\Lambda^n} a_\lambda s_\lambda(t_1, ..., t_n).$$

We wish to show that, as in $R_n^G(W)$, a single Hilbert series determines all the coefficients $\{a_\lambda\}_{\lambda\in\Lambda}$, and that Theorem 1.5 can be applied to this series. Fix an integer t,

and recall that for every polynomial $GL_n(F)$ representation Y the t homogenous sub representation of Y is $Y^{(t)} = \bigoplus_{|\alpha|=t} Y^{\alpha}$. Let us define for every $s = 0, ..., t$ the spaces $Q_s = R_{n,e}^{(t-s)} B_n^{(s)} \subset \left(R_n^G(W)\right)^{(t)}$ where $R_{n,e} = {}^{F\langle x_{1,e},...,x_{n,e}\rangle}/_{Id^G(W)\cap F\langle x_{1,e},...,x_{n,e}\rangle}$, and the spaces $M_p = \sum_{s=p}^{t} Q_s$ for $p = 0, ..t$ (and $M_{t+1} = \{0\}$). It is easy to see that Q_s and M_p are both polynomial $GL_n(F)$ representations for every s and p.

Lemma 3.1.

(1) $\left(R_n^G(W)\right)^{(t)} = M_0$
(2) $\left(R_n^G(W)\right)^{(t)} \cong \bigoplus_{p=0}^{t} {}^{M_p}/_{M_{p+1}}$ *as* $GL_n(F)$ *modules.*

Proof. It is clear that $M_0 \subset \left(R_n^G(W)\right)^{(t)}$. For the other inclusion, note that the free algebra $F\left\langle X_n^G \right\rangle$ is a universal enveloping of the free Lie algebra on X_n^G. Hence, according to Poincare-Birkhoff-Witt theorem, if $l_1, l_2, ...$ is a totally ordered basis of the free Lie algebra, then $F\left\langle X_n^G \right\rangle$ has a basis $\left\{ l_1^{\beta_1} \cdots l_r^{\beta_r} |\ \beta_i \geq 0 \right\}$. In our case we choose $l_i = x_{i,e}$ for $i = 1, ..., n$, and the other basis elements are arranged in an order such that $deg(l_j) \leq deg(l_{j+1})$ for every $j > n$. We conclude that there is a basis K of $F\left\langle X_n^G \right\rangle$ consisting of polynomials of the form $x_{1,e}^{\alpha_1} \cdots x_{n,e}^{\alpha_n} x_{i_1,h_1} \cdots x_{i_r,h_r} [x_{i_{r+1},g_1}, ...] \cdots [..., x_{i_s,g_s}]$ which is in $R_{n,e}^{(|\alpha|)} B_n^{(s)}$ since $h_i \neq e$ for every $i = 1, ..., r$. Moreover, the quotient map is a G-graded homomorphism from $F\left\langle X_n^G \right\rangle^{(t)}$ onto $\left(R_n^G(W)\right)^{(t)}$, thus $\left(R_n^G(W)\right)^{(t)}$ is spanned by the t-homogenous elements of K (modulo the identities).

For the second part recall that every $GL_n(F)$ representation is completely reducible, i.e. $\left(R_n^G(W)\right)^{(t)} = \bigoplus_{i\in I} V_i$ where I is a finite set of indexes and V_i is irreducible $GL_n(F)$ representation for every $i \in I$. Moreover, $Q_s = \bigoplus_{i\in I_s} V_i$ where I_s is a subset of I, and $M_p = \bigoplus_{i\in \cup_{s=p}^{t} I_s} V_i$. Therefore, ${}^{M_p}/_{M_{p+1}} \cong \bigoplus_{i\in J_p} V_i$ where $J_p = \bigcup_{s=p}^{t} I_s - \bigcup_{s=p+1}^{t} I_s$. So to prove the second part we need to show that I is a disjoint union of J_p where $p = 0, ..., t$. The inclusion $\bigcup_{p=0}^{t} J_p \subset I$ is trivial. For the other direction, recall that by part one $I = \bigcup_{s=0}^{t} I_s$, so for every $i \in I$ there is a minimal p such that $i \in I_p$. Hence i belongs to $\bigcup_{s=p}^{t} I_s$ and not to $\bigcup_{s=p+1}^{t} I_s$. By definition, $i \in J_p$, and we conclude that $I = \bigcup_{p=0}^{t} J_p$. To show that the J_p's are disjoint suppose that there is an index i in $J_p \cap J_{p'}$ for $p > p'$. The set J_p is a subset of I_p, and $p \geq p' + 1$, thus $i \in \bigcup_{s=p'+1}^{t} I_s$. Hence, by definition, $i \notin J_{p'}$, which is a contradiction. □

The next step is to find a basis of ${}^{M_p}/_{M_{p+1}}$ for every $p \leq t$. In order to achieve this, we need some preparations:

Lemma 3.2. *Let* $\{y_1, ..., y_k\}$ *be a set of non commutative variables where* $k \geq 2$*, then we have the following identity*

$$y_1 \cdots y_k = y_2 y_1 y_3 \cdots y_k + \sum_{i=0}^{k-2} \sum y_{j_1} \cdots y_{j_i} [y_{l_1}, ..., y_{l_{k-i}}] \tag{3.1}$$

where the inner sum is over some subset of

$$\left\{(j_1,...,j_i,l_1,...,l_{k-i}) \in \mathbb{N}^k \mid \{j_1,...,j_i,l_1,...,l_{k-i}\} = \{1,...,k\}\right\}.$$

Proof. By induction on k. The base of the induction, $k=2$, is true by the definition of the commutator $y_1y_2 = y_2y_1 + [y_1,y_2]$. For the induction step let us assume that the claim is true for k. So, by multiplying the identity 3.1 from the right by y_{k+1} we get:

$$y_1 \cdots y_{k+1} = y_2y_1y_3 \cdots y_{k+1} + \sum_{i=0}^{k-2} \sum y_{j_1} \cdots y_{j_i}[y_{l_1},...,y_{l_{k-i}}]y_{k+1}.$$

We complete the proof by using the commutator definition $zw = wz + [z,w]$ with $z = [y_1,...,y_{l_{k-i}}]$ and $w = y_{k+1}$ for every element in the sum. □

Definition 3.3. We say that $f \in F\left\langle X_n^G \right\rangle$ is *g multi-homogenous of degree* $\gamma = (\gamma_1,...,\gamma_n)$, if substituting the variables $x_{i,g}$ by $t_ix_{i,g}$ in f gives

$$\left(\prod_{i=1}^{n} t_i^{\gamma_i}\right) f.$$

In other words for every $i = 1,..,n$ the variable $x_{i,g}$ appears γ_i times in every monomial of f. We denote the g multi-homogenous degree of polynomial f by $deg_g(f)$.

It is known that for any $g \in G$ if $f = \sum_\gamma f_\gamma$ is a G-graded identity, where f_γ is g multi-homogenous of degree γ, then f_γ is also G-graded identity for every γ. Therefore, the decomposition $B_n^{(p)} = \bigoplus_\gamma \left(B_n^{(p)}\right)_\gamma$ where $\left(B_n^{(p)}\right)_\gamma$ is the subspace of $B_n^{(p)}$ consisting of g multi-homogenous polynomials of degree γ. We may therefore assume that for every $g \in G$ there is a basis $\left\{f_j^{(p)}(x_{l,h}) \mid j = 1,...,dim_F B_n^{(p)}\right\}$ of $B_n^{(p)}$ consisting of g multi-homogenous polynomials. Now we are ready to present the basis of M_p/M_{p+1}.

Proposition 3.4. *For every $p \le t$ let $\left\{f_j^{(p)}(x_{l,h}) \mid j = 1,...,dim_F B_n^{(p)}\right\}$ be a basis of $B_n^{(p)}$ consisting of e multi-homogenous polynomials. Then:*

(1) *For every $1 \le j \le dim_F B_n^{(p)}$ and $\sigma \in S_{t-p}$*

$$x_{i_1,e} \cdots x_{i_{t-p},e} f_j^{(p)}(x_{l,h}) \equiv x_{i_{\sigma(1)},e} \cdots x_{i_{\sigma(t-p)},e} f_j^{(p)}(x_{l,h}) \ (mod\ M_{p+1})$$

(2) *The set*

$$U_p = \left\{x_{1,e}^{\alpha_1} \cdots x_{n,e}^{\alpha_n} f_j^{(p)}(x_{l,h}) \mid |\alpha| = t-p,\ j = 1,...,dim_F B_n^{(p)}\right\}$$

is a basis of M_p/M_{p+1}.

Proof. To prove the first part it is enough to show that for every $1 \le r \le t-p-1$

$$x_{i_1,e} \cdots x_{i_{t-p},e} f_j^{(p)}(x_{l,h}) \equiv x_{i_1,e} \cdots x_{r+1,e} x_{r,e} \cdots x_{i_{t-p},e} f_j^{(p)}(x_{l,h}) \pmod{M_{p+1}}.$$

Recall that $f_j^{(p)}$ is a linear combination of polynomials of the form $x_{i_1,h_1} \cdots x_{i_q,h_q}[x_{l_1,g_1},...,][...,x_{l_{p-q},g_{p-q}}]$ where $h_i \ne e$. Hence, by using Lemma 3.2 on

$x_{r,e}\cdots x_{i_{t-p},e}x_{i_1,h_1}\cdots x_{i_q,h_q}$ for every such monomial, we obtain

$$x_{i_1,e}\cdots x_{i_{t-p},e}f_j^{(p)}(x_{l,h}) = x_{i_1,e}\cdots x_{i_{r+1},e}x_{i_r,e}\cdots x_{i_{t-p},e}f_j^{(p)}(x_{l,h}) + P(x_{l,h})$$

where $P(x_{l,h})$ is a linear combination of polynomials of the form $x_{j_1e},...,x_{j_{t-q-k},e}x_{i_1,h_1}\cdots x_{i_q,h_q}[x_{l_1,g_1},...,][...,x_{l_k,g_k}]$ with $k+q>p$. Thus $P(x_{l,h})\in M_{p+1}$, and the first part is proven.

Obviously U_p spans ${}^{M_p}/_{M_{p+1}}$. It remains to be show that U_p is linearly independent, so let

$$T=\sum_{j,\alpha}\beta_{j,\alpha}x_{1,e}^{\alpha_1}\cdots x_{n,e}^{\alpha_n}f_j^{(p)}(x_{l,h})$$

be a nontrivial linear combination of the elements in U_p. Assume by contradiction that T is an identity of W. Therefore T_γ, the e multi-homogenous component of degree γ of T, is also an identity for every $\gamma\in\mathbb{N}^n$. We choose γ such that T_γ is a nontrivial linear combination, and write

$$T_\gamma=\sum_j\sum_{\alpha=\gamma-deg_e\left(f_j^{(p)}\right)}\beta_{j,\alpha}x_{1,e}^{\alpha_1}\cdots x_{n,e}^{\alpha_n}f_j^{(p)}$$

where the first sum is over every $1\le j\le dim_F B_n^{(p)}$ such that $deg_e\left(f_j^{(p)}\right)_i\le\gamma_i$ for every $i=1,...,n$ (note that every index j is appearing at most once in the sum).

Now, let substitute $x_{i,e}$ by $x_{i,e}+1$ in T_γ and denote the obtained identity by S. Notice that since $[y+1,z]=[y,z]$ the polynomials $f_j^{(p)}$ are stay the same under this substitution (in the part which is not commutator products in $f_j^{(p)}$ there are no e-homogenous variables). Therefore, the p - homogenous component of S is

$$\sum_j\beta_{j,\alpha}f_j^{(p)}$$

which is also an identity of W. But $f_j^{(p)}$ are linear independent (modulo the identities) and not all the $\beta_{j,\alpha}$ are zero, and this is a contradiction . □

Corollary 3.5. .

(1) $R_n^G(W)\cong F[z_1,...,z_n]\otimes B_n$ *as* $GL_n(F)$ *representation.*
(2) $c_n^G(W)=\sum_{s=0}^n\binom{n}{s}\delta_s$ *where* $\delta_s=dim_F B_s^{(1^s)}$.

Proof. By 3.4 the mapping from ${}^{M_p}/_{M_{p+1}}$ to $F[z_1,...,z_n]^{(t-p)}\otimes B_n^{(p)}$ taking $\left(\prod_{i=1}^n x_{i,e}^{\alpha_i}\right)f_j^{(p)}$ to $\left(\prod_{i=1}^n z_i^{\alpha_i}\right)\otimes f_j^{(p)}$ is a $GL_n(F)$ representation isomorphism, thus by Lemma 3.1 $R_n^G(W)^{(t)}\cong\left(F[z_1,...,z_n]\otimes B_n\right)^{(t)}$ for every t.

For the proof of the second part, note that

$$C_n^G(W)\cong\left(F[z_1,...,z_n]\otimes B_n\right)^{(1^n)}=\bigoplus_{\beta\in\{0,1\}^n}F[z_1,...,z_n]^{(1^n)-\beta}\otimes B_n^\beta$$

The dimension of $F[z_1,...,z_n]^{(1^n)-\beta}$ is 1 for every $\beta \in \{0,1\}^n$. Furthermore, it is easy to see that $B_n^\beta \cong B_{|\beta|}^{(1^{|\beta|})}$ for every $\beta \in \{0,1\}^n$, and there are $\binom{n}{s}$ vectors $\beta \in \{0,1\}^n$ such that $|\beta| = s$. Thus $dim_F C_n^G(W) = c_n^G(W) = \sum_{s=0}^n \binom{n}{s}\delta_s$. □

With this corollary we can write

$$H_{R_n^G(W)} = H_{F[z_1,...,z_n]} \cdot H_{B_n}.$$

Recall that $H_{R_n^G(W)}(t_1,...,t_n) = \sum_\lambda m_\lambda s_\lambda(t_1,...,t_n)$ and $H_{B_n}(t_1,...,t_n) = \sum_\lambda a_\lambda s_\lambda(t_1,...,t_n)$. Moreover, the Hilbert series of $F[z_1,...,z_n]$ is known to be

$$\sum_{\alpha\in\mathbb{N}^n} t_1^{\alpha_1}\cdots t_n^{\alpha_n} = \prod_{i=1}^n (1-t_i)^{-1} = \sum_{k\in\mathbb{N}} s_{(k)}(t_1,...,t_n), \tag{3.2}$$

thus we have the identity

$$\sum_\lambda m_\lambda s_\lambda = \sum_{\mu\in\Lambda}\sum_{k\in\mathbb{N}} a_\mu s_\mu s_{(k)} \tag{3.3}$$

This identity gives us the following arithmetic connection between m_λ and a_μ.

Proposition 3.6. *For every partition λ*

$$m_\lambda = \sum_{\mu\in L_\lambda} a_\mu$$

where L_λ is the set of all partition μ such that $\lambda_{i+1} \le \mu_i \le \lambda_i$ for every $i = 1,...,h(\mu)$.

Proof. By Pieri's formula (see [M]) $s_\mu s_{(k)} = \sum s_\nu$ where the sum is over all partitions ν obtained from μ by adding k boxes to its Young's diagram, no two of them in the same column. We have to show that s_λ appearing in the sum on right hand side of $s_\mu s_{(|\lambda|-|\mu|)} = \sum s_\nu$ if and only if $\lambda_{i+1} \le \mu_i \le \lambda_i$ for every i.

Suppose that s_λ appearing in the sum on the right hand side of $s_\mu s_{(|\lambda|-|\mu|)} = \sum s_\nu$. It is obvious that $\mu_i \le \lambda_i$ for every i, since we are adding boxes to μ in order to get λ. Let us assume that there is an index i such that $\mu_i < \lambda_{i+1} \le \lambda_i$. In the$\mu_i + 1$ column, then, we necessarily added two boxes which is a contradiction. (see Figure 3.1)



FIGURE 3.1.

For the other direction, let λ and μ be partitions such that $\lambda_{i+1} \le \mu_i \le \lambda_i$ for every i. Suppose that in the j's column we added two boxes in the k 's and i's row where $k > i$. This implies that $\mu_i < j \le \lambda_k \le \lambda_{i+1}$ and this is a contradiction. □

The conclusion of the above proposition is that the coefficients a_λ inherit two properties from the coefficients m_λ. Here is the first one.

Corollary 3.7. *There exists an integer q such that $a_\lambda = 0$ if $\lambda \notin \Lambda^q$.*

Proof. Recall from Section 1 that $m_\lambda = 0$ if $\lambda \notin \Lambda^{dim_F(A_W)}$ where A_W is the finite dimensional algebra with the same G-graded identities as W.

Let μ be a partition such that $h(\mu) > dim_F(A_W) - 1$ (i.e. $\mu \notin \Lambda^{dim_F A_W - 1}$), and define $\lambda = (\mu_1, ..., \mu_{h(\mu)}, 1)$. Since $h(\lambda) = h(\mu) + 1 > dim_F(A_W)$ we know that $m_\lambda = \sum_{\nu \in L_\lambda} a_\nu = 0$, so $a_\nu = 0$ for any $\nu \in L_\lambda$. Moreover, it is easy to see that $\lambda_{i+1} \leq \mu_i \leq \lambda_i$ for every i thus $\mu \in L_\lambda$. □

By this corollary, and since $H_{B_n}(t_1, ..., t_n) = \sum_{\lambda \in \Lambda^n} a_\lambda s_\lambda(t_1, ..., t_n)$ for every n, we conclude that there is a single Hilbert series (here it is $H_{B_q}(t_1, ..., t_q)$) which determines all the coefficients a_λ. The second corollary of 3.6 is that, as in $R_n^G(W)$ (see Section 1), the series $H_{B_q}(t_1, .., t_q)$ satisfies the first assumption of Theorem 1.5.

Corollary 3.8. *Let $H_{B_q}(t_1, ..., t_q) = \sum_\lambda a_\lambda s_\lambda(t_1, ..., t_q)$. Then there is an integer r such that $a_\lambda = 0$ for every partition λ satisfying $\lambda_{r+1} \geq M$ where M is a constant depend only on r.*

Moreover the integer $r = exp^G(W) - 1$ is the minimal satisfying this condition.

Proof. From [Go] there exists an integer M such that $m_\lambda = 0$ for every partition λ satisfies $\lambda_{l+1} \geq M$ where $l = exp^G(W)$. Suppose that μ is a partition such that $\mu_l \geq M$. We define a partition $\lambda = (\mu_1, ..., \mu_l, \mu_l, \mu_{l+2}, ..., \mu_{h(\mu)})$. Obviously, $\lambda_{l+1} = \mu_l \geq M$, so $m_\lambda = \sum_{\nu \in L_\lambda} a_\nu = 0$, so $a_\nu = 0$ for any $\nu \in L_\lambda$. Moreover, one can check that $\lambda_{i+1} \leq \mu_i \leq \lambda_i$ for every i so $\mu \in L_\lambda$.

To show the minimality of $l - 1$, we have to show that for every integer p there is a partition μ with $\mu_{l-1} \geq p$ such that $a_\mu \neq 0$. We know from [Go] that for every integer p there is a partition λ with $\lambda_l \geq p$ such that $m_\lambda = \sum_{\nu \in L_\lambda} a_\nu \neq 0$. Hence, there is $\mu \in L_\lambda$, which satisfies $\mu_{l-1} \geq \lambda_l \geq p$, such that $a_\mu \neq 0$. □

The Hilbert series H_{B_q} is also nice rational because $H_{R_q^G(W)}$ is a nice rational function, and since 3.2 imply that

$$H_{B_q}(t_1, ..., t_q) = \prod_{i=1}^{q} (1 - t_i) H_{R_q^G(W)}(t_1, ..., t_q).$$

We get that the conclusion of Theorem 1.5 holds for the series $\{\delta_s\}_{s \in \mathbb{N}}$, and together with 3.5

$$c_n^G(W) \sim \sum_{m=0}^{d-1} \sum_{\substack{0 \leq s \leq n \\ s = m \, (mod \, d)}} \binom{n}{s} \alpha_m s^{\beta_m} (l-1)^s$$

where $\beta_m \in \frac{1}{2}\mathbb{Z}$ for every m. To estimate this sum, let ω be a primitive d^{th} root of unit and note that $\sum_{t=0}^{d-1} \omega^{(s-m)t}$ is zero if $s \neq m\,(mod\,d)$, and is d if $s = m\,(mod\,d)$. Thus,

$$c_n^G(W) \sim \sum_{m=0}^{d-1} \sum_{\substack{0 \leq s \leq n \\ s = m\,(mod\,d)}} \binom{n}{s} \alpha_m s^{\beta_m} (l-1)^s$$

$$= \frac{1}{d} \sum_{m=0}^{d-1} \sum_{s=0}^{n} \sum_{t=0}^{d-1} \omega^{(s-m)t} \binom{n}{s} \alpha_m s^{\beta_m} (l-1)^s$$

$$= \sum_{m=0}^{d-1} \sum_{t=0}^{d-1} \frac{\alpha_m}{d} \omega^{-mt} \sum_{s=0}^{n} \binom{n}{s} s^{\beta_m} \left(\omega^t (l-1)\right)^s$$

By lemma 1.1 in [BeR], for every m and t the expression $\sum_{s=0}^{n} \binom{n}{s} s^{\beta_m} \left(\omega^t (l-1)\right)^s$ asymptotically equal to:

$$\mu_{m,t} n^{\beta_m} \left(\omega^t (l-1) + 1\right)^n$$

where $\mu_{m,t}$ is a constant which does not depend on n. Moreover, the absolute value of the expression $\omega^t(l-1)+1$ is maximal when $t = 0$, so

$$c_n^G(W) \sim \alpha n^\beta l^n$$

where $\beta = max(\beta_m) \in \frac{1}{2}\mathbb{Z}$. And theorem **B** is proven.

References

[AB1] E. Aljadeff and A. Kanel-Belov, Hilbert series of PI relatively free G-graded algebras are rational functions. Bull. Lond. Math. Soc. 44 (2012), no. 3, 520–532.

[AB2] E. Aljadeff and A. Kanel-Belov, 'Representability and Specht problem for G-graded algebras', Adv. Math. 225 (2010) 2391–2428

[AG] E. Aljadeff and A. Giambruno, Multialternating graded polynomials and growth of polynomial identities, to appear in Proc. Amer. Math. Soc.

[AGL] E. Aljadeff, A. Giambruno and D. La Mattina, Graded polynomial identities and exponential growth, J. Reine Angew. Math. 650 (2011), 83–100.

[BeR] W.Beckner and A.Regev, Asymptotic estimates using probability. Adv. Math. 138 (1998), no. 1, 1–14.

[BRo] A. Kanel-Belov and L.H. Rowen, Computational aspects of polynomial identities. Research Notes in Mathematics, 9. A K Peters, Ltd., Wellesley, MA, 2005.

[BR] A.Berele, and A.Regev, Asymptotic behaviour of codimensions of p. i. algebras satisfying Capelli identities. Trans. Amer. Math. Soc. 360 (2008), no. 10, 5155–5172.

[Bru] Bruce .E. Sagan, The Symmetric Group: Representations, Combinatorial Algorithms and Symmetric Functions, Section 4.4, 155-157

[D] V. Drensky, Codimensions of T-ideals and Hilbert series of relatively free algebras, J. of Alg. 89 (1984), 178–223. MR765766 (86b:16010)

[GL] A. Giambruno and D. La Mattina, Graded polynomial identities and codimensions: computing the exponential growth. Adv. Math. 225 (2010), no. 2, 859–881.

[GZ1] A. Giambruno and M. V. Zaicev, On codimension growth of finitely generated associative algebras, Adv. Math. 140 (1998), 145–155.

[GZ2] Giambruno, Antonio; Zaicev, Mikhail, Polynomial identities and asymptotic methods. Mathematical Surveys and Monographs, 122. American Mathematical Society, Providence, RI, 2005.

[GZ3] A.Giambruno; M.V.Zaicev, Growth of polynomial identities: Is the sequence of codimensions eventually nondecreasing? Bull. Lond. Math. Soc. 46 (2014), no. 4, 771–778.

[Go] A. S.Gordienko, Amitsur's conjecture for associative algebras with a generalized Hopf action. J. Pure Appl. Algebra 217 (2013), no. 8, 1395–1411.

[M] I. G.Macdonald, (1995), Symmetric functions and Hall polynomials, Oxford Mathematical Monographs (2nd ed.), The Clarendon Press Oxford University Press

[St] R. Stanley, Combinatorics and Commutative Algebra, Birkhäuser, Boston, 1983.

[Stu] B. Sturmfels, On vector partition functions, J. Combin. Theory Ser. A 72 (1995) 302–308.

[Sv] I.Sviridova, Identities of pi-algebras graded by a finite abelian group. (English summary) Comm. Algebra 39 (2011), no. 9, 3462–3490.

Department of Mathematics, Technion - Israel Institute of Technology, Haifa 32000, Israel.

E-mail address: `yuvalshp 'at' tx.technion.ac.il`